

$X(3872)$ in Effective Field Theory

S. Fleming* and T. Mehen†

*Physics Department, University of Arizona, Tucson, AZ 85721

†Physics Department, Duke University, Durham NC, 27708

Abstract.

If the $X(3872)$ resonance is a shallow boundstate of the charm mesons $D^0\bar{D}^{*0}$ and $D^{*0}\bar{D}^0$, it can be described by an effective theory of nonrelativistic D mesons coupled to nonrelativistic pions (X-EFT). In this talk, I give a brief overview of the $X(3872)$, followed by a short review of X-EFT. I end my talk with results from calculations of the next-to-leading-order correction to the partial decay width $\Gamma[X \rightarrow D^0\bar{D}^0\pi^0]$, and the decay of $X(3872)$ to P -wave quarkonia.

Keywords: Molecular Charm

PACS: 12.39.Fe, 12.39.Hg, 14.40.Lb

INTRODUCTION

The $X(3872)$ [1, 2, 3] is a novel charmonium state that lies very close to the $D^0\bar{D}^{*0}$ threshold. The closeness of the $X(3872)$ to this threshold ($-0.6 \pm 0.6\text{MeV}$) has prompted numerous authors to suggest that it is a $C = 1$ molecular bound state of $D^0\bar{D}^{*0} + \bar{D}^0D^{*0}$, for reviews see Refs. [4, 5]. The strongest evidence for the molecular hypothesis is the large ratio of the branching fractions for the $X(3872)$ decays to J/ψ plus two or three pions [6]

$$\frac{\text{Br}[X(3872) \rightarrow J/\psi \pi^+ \pi^- \pi^0]}{\text{Br}[X(3872) \rightarrow J/\psi \pi^+ \pi^-]} = 1.0 \pm 0.4 \pm 0.3. \quad (1)$$

The final states have opposite G-parity which implies that the $X(3872)$ does not have definite isospin. Measurements of the invariant mass distributions of the pions indicate that these decays proceed through $J/\psi \rho$ for the $J/\psi \pi^+ \pi^-$ final state, and through $J/\psi \omega$ for the $J/\psi \pi^+ \pi^- \pi^0$ final state, suggesting that the $X(3872)$ couples to $I = 0$ and $I = 1$ channels with roughly equal strength. This is not possible for a conventional $c\bar{c}$ state. The observed branching ratio can be understood if the $X(3872)$ is comprised primarily of neutral D mesons, which is expected since the charged $D-D^*$ threshold is far (8.5 MeV) above the $X(3872)$ mass. There is, however, a caveat, as pointed out in Refs. [7, 8] the ratio above could also be due to the kinematic suppression of the $J/\psi \omega$ mode, which leaves an opening for a charmonium interpretation of the $X(3872)$. Another possibility, suggested in Ref. [9], is that there are actually two nearly degenerate states with opposite G-parity.

X-EFT

If the $X(3872)$ is a molecular state then the large distance behavior of the $D\bar{D}^*$ can be described by X-EFT [10, 11], an effective field theory which will be described below. The upper bound on the typical momentum of the D and $\bar{D}^*$ in the bound state is $\gamma \equiv (2\mu_{D\bar{D}^*}E_X)^{1/2} \leq 48\text{MeV}$, where $\mu_{D\bar{D}^*}$ is the reduced mass of the D^0 and $\bar{D}^{*0}$. For this binding momentum the typical velocity of the D and D^* is approximately $v_D \simeq (E_X/2\mu_{D\bar{D}^*})^{1/2} \leq 0.02$, and both the D and D^* are non-relativistic.

The pion degrees of freedom are also treated non-relativistically. The maximum energy of the pion emitted in the decay $X \rightarrow D^0\bar{D}^0\pi^0$ is

$$E_\pi = \frac{m_X^2 - 4m_D^2 + m_\pi^2}{2m_X} = 142\text{MeV}, \quad (2)$$

which is just 7 MeV above the π^0 mass at 134.98 MeV. The maximum pion momentum is approximately 44 MeV, which is comparable to both the typical D -meson momentum, $p_D \sim \gamma \leq 48\text{MeV}$, and the momentum scale appearing in the pion-exchange graph: $\mu \simeq 45\text{MeV}$. Since the velocity of the pions is $v_\pi = p_\pi/m_\pi \leq 0.34$, a non-relativistic treatment of the pion fields is valid.

The effective Lagrangian includes the charm mesons, the anti-charm mesons, and the pion fields. We denote the fields that annihilate the D^{*0} , $\bar{D}^{*0}$, D^0 , $\bar{D}^0$, and π^0 as $\mathbf{D}$, $\bar{\mathbf{D}}$, D , $\bar{D}$, and π , respectively. To the order we are working we will not need diagrams with charged pions and charged D mesons so these are neglected in what follows. We construct an effective Lagrangian that is relevant for low-energy S -wave $D\bar{D}^*$ scattering, where the initial and the final states are the $C = +$ superposition of $D^0\bar{D}^{*0}$ and $D^{*0}\bar{D}^0$:

$$|D\bar{D}^*\rangle \equiv \frac{1}{\sqrt{2}} [|D^0\bar{D}^{*0}\rangle + |D^{*0}\bar{D}^0\rangle]. \quad (3)$$

An interpolating field with these quantum numbers will be used to calculate the properties of the $X(3872)$. We integrate out all momentum scales much larger than the momentum scale set by $p_D \sim p_\pi \sim \mu$. For D mesons this corresponds to kinetic energy $\leq 1\text{MeV}$, for pions the kinetic energy is $\leq 7\text{MeV}$. The hyperfine splitting Δ and m_π should be treated as large compared to the typical energy scale in the theory. The effective X-EFT Lagrangian is

$$\begin{aligned} \mathcal{L} = & \mathbf{D}^\dagger \left(i\partial_0 + \frac{\vec{\nabla}^2}{2m_{D^*}} \right) \mathbf{D} + D^\dagger \left(i\partial_0 + \frac{\vec{\nabla}^2}{2m_D} \right) D + \bar{\mathbf{D}}^\dagger \left(i\partial_0 + \frac{\vec{\nabla}^2}{2m_{D^*}} \right) \bar{\mathbf{D}} \\ & + \bar{D}^\dagger \left(i\partial_0 + \frac{\vec{\nabla}^2}{2m_D} \right) \bar{D} + \pi^\dagger \left(i\partial_0 + \frac{\vec{\nabla}^2}{2m_\pi} + \delta \right) \pi \\ & + \left(\frac{g}{\sqrt{2}f_\pi} \right) \frac{1}{\sqrt{2}m_\pi} \left(D\mathbf{D}^\dagger \cdot \vec{\nabla}\pi + \bar{D}^\dagger\bar{\mathbf{D}} \cdot \vec{\nabla}\pi^\dagger \right) + \text{h.c.} \\ & - \frac{C_0}{2} (\bar{\mathbf{D}}D + \mathbf{D}\bar{\mathbf{D}})^\dagger \cdot (\bar{\mathbf{D}}D + \mathbf{D}\bar{\mathbf{D}}) \end{aligned}$$

$$\begin{aligned}
& + \frac{C_2}{16} (\bar{\mathbf{D}}\mathbf{D} + \mathbf{D}\bar{\mathbf{D}})^\dagger \cdot (\bar{\mathbf{D}}(\nabla)^2\mathbf{D} + \mathbf{D}(\nabla)^2\bar{\mathbf{D}}) + h.c. \\
& + \frac{B_1}{\sqrt{2}} \frac{1}{\sqrt{2m_\pi}} (\bar{\mathbf{D}}\mathbf{D} + \mathbf{D}\bar{\mathbf{D}})^\dagger \cdot D\bar{\mathbf{D}}\vec{\nabla}\pi + h.c. + \dots,
\end{aligned} \tag{4}$$

where $\delta = \Delta - m_\pi \simeq 7 \text{ MeV}$, ∇ is the operator $\vec{\nabla}$ operating to the right minus $\vec{\nabla}$ operating to the left, $\mu^2 = \Delta^2 - m_\pi^2 \approx 2m_\pi\delta$, and $\dots$ denotes higher-order interactions. The pion decay constant is $f_\pi = 132 \text{ MeV}$ with our choice of normalization. Additional terms that couple the $X(3872)$ to charmonium states can be added to the lagrangian above. For example, the coupling of the χ_{cJ} to heavy mesons is given by

$$\mathcal{L}_\chi = i\frac{g_1}{2} \text{Tr}[\chi^{\dagger i} H_a \sigma^i \bar{H}_a] + \frac{c_1}{2} \text{Tr}[\chi^{\dagger i} H_a \sigma^j \bar{H}_b] \epsilon_{ijk} A_{ab}^k + h.c.. \tag{5}$$

Note the appearance of additional unknown couplings: g_1 and c_1 .

RESULTS

Using the lagrangian given at the end of the last section it is straightforward to calculate the the next-to-leading-order correction to the partial decay width $\Gamma[X \rightarrow D^0 \bar{D}^0 \pi^0]$, as well as the decays $X(3872) \rightarrow \chi_{cJ}(\pi^0, \pi\pi)$. In Fig. 1 we present the LO and NLO predictions for the decay rate for $X \rightarrow D^0 \bar{D}^0 \pi^0$. Note, the LO decays $X(3872) \rightarrow D^0 \bar{D}^0 \pi^0$ was first calculated in Ref. [12], in addition the decay $X(3872) \rightarrow D^0 \bar{D}^0 \gamma$ was calculated in Ref.[13]. At NLO the decay rate depends on two new undetermined parameters, r and η . In order to estimate their effect we vary these parameters within a reasonable range.

Using these results we can calculate the partial rates for $\Gamma[X(3872) \rightarrow \chi_{cJ} \pi^0]$. At LO ($c_1 = 0$) we find (denoting $\Gamma[X(3872) \rightarrow \chi_{cJ} \pi^0] \equiv \Gamma_J$)

$$\Gamma_0 : \Gamma_1 : \Gamma_2 :: 4.76 : 1.57 : 1. \tag{6}$$

The energy dependence from the virtual pion propagators makes a significant change in the predictions for the relative sizes of the partial rates. The NLO predictions depend on the ratio c_1/g_1 which is undetermined. The ratio c_1/g_1 has dimensions of inverse mass, when c_1/g_1 is varied from $(100 \text{ MeV})^{-1}$ to $(1000 \text{ MeV})^{-1}$, we find:

$$\begin{aligned}
\Gamma_0 : \Gamma_1 : \Gamma_2 &:: 3.01 : 1.06 : 1 & c_1/g_1 &= (100 \text{ MeV})^{-1} \\
\Gamma_0 : \Gamma_1 : \Gamma_2 &:: 3.49 : 1.20 : 1 & c_1/g_1 &= (300 \text{ MeV})^{-1} \\
\Gamma_0 : \Gamma_1 : \Gamma_2 &:: 3.76 : 1.28 : 1 & c_1/g_1 &= (500 \text{ MeV})^{-1} \\
\Gamma_0 : \Gamma_1 : \Gamma_2 &:: 4.11 : 1.38 : 1 & c_1/g_1 &= (1000 \text{ MeV})^{-1}.
\end{aligned} \tag{7}$$

For the largest values of c_1/g_1 , we obtain predictions for the relative sizes of the partial rates similar to those found in Ref. [14]. As c_1/g_1 decreases, the predictions tend to those appearing in Eq. (6). Experimental measurement of the relative sizes of the partial rates for decays to $\chi_{c,J} \pi^0$ can be used to determine c_1/g_1 .




FIGURE 1. Decay rate for $X \rightarrow D^0 \bar{D}^0 \pi^0$ as a function of E_X . We use $g = 0.6$. The central solid line corresponds to the LO prediction. The band is the result of the NLO calculation when the parameters r_0 and η are varied in the ranges $0 \leq r_0 \leq (100 \text{ MeV})^{-1}$ and $-1 \leq \eta \leq 1$.

ACKNOWLEDGMENTS

This work was supported in part by the Director, Office of Science, Office of Nuclear Physics, of the U.S. Department of Energy under grant number DE-FG02-06ER41449.

REFERENCES

1. S. K. Choi *et al.* [Belle Collaboration], Phys. Rev. Lett. **91**, 262001 (2003) [arXiv:hep-ex/0309032].
2. D. Acosta *et al.* [CDF II Collaboration], Phys. Rev. Lett. **93**, 072001 (2004) [arXiv:hep-ex/0312021].
3. V. M. Abazov *et al.* [D0 Collaboration], Phys. Rev. Lett. **93**, 162002 (2004) [arXiv:hep-ex/0405004].
4. M. B. Voloshin, in *Proceedings of 4th Flavor Physics and CP Violation Conference (FPCP 2006), Vancouver, British Columbia, Canada, 9-12 Apr 2006*, pp 014 [arXiv:hep-ph/0605063].
5. S. Godfrey and S. L. Olsen, arXiv:0801.3867 [hep-ph].
6. K. Abe *et al.* [Belle Collaboration], arXiv:hep-ex/0505037.
7. M. Suzuki, Phys. Rev. D **72**, 114013 (2005) [arXiv:hep-ph/0508258].
8. F. De Fazio, Phys. Rev. D **79**, 054015 (2009) [arXiv:0812.0716 [hep-ph]].
9. D. Gamermann and E. Oset, Eur. Phys. J. A **33**, 119 (2007) [arXiv:0704.2314 [hep-ph]].
10. S. Fleming, M. Kusunoki, T. Mehen and U. van Kolck, Phys. Rev. D **76**, 034006 (2007) [arXiv:hep-ph/0703168].
11. S. Fleming and T. Mehen, Phys. Rev. D **78**, 094019 (2008) [arXiv:0807.2674 [hep-ph]].
12. M. B. Voloshin, Phys. Lett. B **579**, 316 (2004) [arXiv:hep-ph/0309307].
13. M. B. Voloshin, Int. J. Mod. Phys. A **21**, 1239 (2006) [arXiv:hep-ph/0509192].
14. S. Dubynskiy and M. B. Voloshin, Phys. Rev. D **77**, 014013 (2008) [arXiv:0709.4474 [hep-ph]].